THRIVING



Helping Adults with Intellectual and Developmental Disabilities Heal and Thrive Following Trauma

PARTICIPANT MANUAL

ACKNOWLEDGEMENTS

This project would not have been made possible without the support and hard work of many people and organizations. These materials have been adapted with permission from "The Road to Recovery Toolkit: Supporting Children with IDD Who Have Experienced Trauma" developed by the National Center for Child Traumatic Stress and the National Child Traumatic Stress Network and a national selection of experts in trauma and IDD. The toolkit is an excellent resource for providers of children with IDD.

This project would not have been possible without the funding of the WITH Foundation. The WITH Foundation works to promote comprehensive healthcare for adults with developmental disabilities that meets their unique and fundamental needs.

The chair would like to extend a heartfelt thank you to the Advisory Board:

Chair:

Ruth Brown, PhD Virginia Commonwealth University Richmond.VA

Ananda Amstadter, PhD Virginia Commonwealth University Richmond, VA

Patricia Hagood Richmond,VA

Colleen Horton Hogg Foundation Houston,TX

Adam Kaul, MD Regional Education Assessment Crisis Services and Habilitation Richmond.VA Susan Ko, PhD Science 37 Los Angeles, CA

Andrea Konig, PhD Bon Secours St. Mary's Hospital Richmond, VA

Michelle (Shell) Schwartz, MA SafePlace Austin, TX

Michael Southam-Gerow, PhD Virginia Commonwealth University Richmond, VA

ACKNOWLEDGEMENTS

The Chair wishes to extend special thanks to:

- The WITH Foundation for their generous funding and support of this project.
- The National Child Traumatic Stress Network for their permission to adapt the Road to Recovery Toolkit.
- The Partnership for People with Disabilities at Virginia Commonwealth University for their support, piloting, and valuable feedback.
- Wadah Al Mulhim for initial design work. Arden Hajaligholi for sound editing of the online workshop. Christian Manu and Michael Matthews for web support. Allison D'Aguilar and Zack Moore for editing.

Recommended Citation

Brown, R. C., Amstadter, A. B., Horton, C., Kaul, A., Ko, S., Konig, A., Schwartz, M., Southam-Gerow, M. (2021). *Thriving: Helping Adults with Intellectual and Developmental Disabilities Heal and Thrive Following Trauma: Participant Manual.* Richmond, VA. Virginia Commonwealth University.

Copyright © 2021 Ruth Brown-Ennis and Virginia Commonwealth University.

INTRODUCTION

Welcome to Thriving: Helping Adults with Intellectual and Developmental Disabilities Heal and Thrive Following Trauma. This workshop is an introduction to trauma-informed concepts and resources. Trauma-informed services are those that focus on awareness of the impacts of trauma and practices designed to provide safety and empowerment to those who have experienced trauma.

Trauma-informed care is a universal intervention that benefits everyone – organizations, service providers, and people supported. Depending on your role, you may wish to further your training to be able to provide more trauma-sensitive or trauma-specific services. This training will provide some recommendations for resources to further your training.

This participant manual is part of the Thriving Toolkit, which also consists of a Facilitator's Guide, Slidekit, and Supplemental Materials. Together, they are designed to teach basic knowledge, skills and values about working with adults with intellectual and developmental disability (IDD) who have had traumatic experiences, and how to use this knowledge to support the person's safety, well-being, happiness, and recovery through trauma-informed practice.

These materials have been adapted with permission from "The Road to Recovery Toolkit: Supporting Children with IDD Who Have Experienced Trauma" developed by the National Center for Child Traumatic Stress and the National Child Traumatic Stress Network and a national selection of experts in trauma and IDD. The Thriving Toolkit provides up-to-date tools and resources that are developmentally appropriate for adults with IDD.

The toolkit was designed to support dissemination and implementation of trauma-informed care by providing complete easy-to-use materials so that anyone can provide the Thriving training in their own community and/or organization. The complete toolkit can be downloaded from the Thriving website. We hope that if you find the Thriving training valuable that you will consider hosting a Thriving workshop of your own.

The Thriving website also hosts the online course. The online course was designed to help those who are unable to attend a Thriving workshop due to time or distance. The online training does not contain the interactive activities which are a valuable part of the Thriving training. However, it still contains critical information and resources to support resilience and recovery of people with IDD who have experienced trauma. It is also a useful resource in preparing to provide the Thriving workshop yourself.

Organization of this course

This course contains six modules. Each module contains a set of learning objectives and are built around 10 Essential Messages. These Essential Messages are the key take-home messages of the training. Each module also includes a number of individual and group activities to help you apply what you are learning to your own role and context. If you are attending this course with others from your own or other organizations, we hope you take this opportunity to share your experience and learn from the experiences of others. At the end of each module you will complete and Action Planning activity where you will identify actionable steps you can take to apply the trauma-informed essential messages to your own work. At the end of the training, you will create a Personal Trauma-Informed Action Plan where you will identify the top 3 action steps that you can commit to implement in your work.

CONTENTS

Acknowledgements	I
Introduction	3
Essential Messages	7
Learning Objectives	8
Module One: Setting the Stage	10
Making The Connection: Why Am I Doing This Work?	21
Action Planning Essential Messages 1 & 2	23
Module Two: Development, IDD, and Trauma	25
Living With IDD	33
<u>Case Vignettes</u>	38
Putting It into Practice Case Vignettes: Trauma, IDD & Development	43
Action Planning Essential Messages 3 & 4	45
Module Three: Traumatic Stress Responses in People with IDD	46
Putting It into Practice Case Vignettes: Responses to Traumatic Experiences	57
12 Core Concepts	58
Action Planning Essential Message 5	60
Module Four:Well-Being and Resilience	61
Local Resources	68
Action Planning Essential Messages 6 & 7	74
Module Five: IDD- and Trauma-Informed Services and Treatment	75
Putting It into Practice: Fish Bowl	85
Putting It into Practice: Case Vignettes: Accessing Services	90

Protective Factors Checklist	91
Action Planning Essential Messages 8 & 9	93
Module Six: Provider Self-Care	95
Stress Warning Signs	98
Self-Care Options	100
Balancing Your Self-Care	102
Action Planning Essential Message 10	107
Personal Trauma-Informed Practice Action Plan	108
References	110

ESSENTIAL MESSAGES

- Know that there's hope; recovery from traumatic experiences is possible.
 (Module 1)
- 2. Recognize that an adult with IDD may have had a traumatic experience(s), which can have profound effects. (Module 1)
- 3. Recognize an adult's developmental level and how IDD and traumatic experiences are affecting his/her functioning. (Module 2)
- 4. Utilize a developmental lens when making meaning of an adult's traumatic experiences and responses. (Module 2)
- 5. Recognize that in the aftermath of trauma, understanding traumatic stress responses is the first step in helping an adult regain their sense of safety, value and quality of life. (Module 3)
- 6. Utilize an IDD- and trauma-informed adult-centered approach to support both the adult and the family. (Module 4)
- 7. Help parents/caregivers, and other professionals in the adult's life, strengthen protective factors. (Module 4)
- 8. Partner with agencies and systems to ensure earlier and more sustained access to services. (Module 5)
- 9. Ensure that trauma-informed adult-centered services, treatments and systems drive the recovery plan. (Module 5)
- 10. Practice ongoing self-care in order to increase effectiveness in delivering high quality support, services and treatment. (Module 6)

LEARNING OBJECTIVES

Mod	ule I: Setting the Stage
	Define Trauma-Informed Care
	Realize that people with IDD are at-risk for traumatic experiences.
	Articulate that recover from traumatic experiences is possible.
	Establish the important role of parents/caregivers and the support network on the road to recovery.
	Describe frameworks that will be used to highlight key concepts throughout the training.
Mod	ule 2: Development, IDD & Trauma
	Define IDD and various types of disabilities.
	, ,
	Describe how IDD and traumatic experiences may disrupt typical development.
	Identify the areas in which a person's functioning is affected by IDD & subsequent adaptations that must be made.
Mod	ule 3:Traumatic Stress Responses
in Pe	ople with IDD
	Identify the types of events that can cause traumatic stress.
	Define traumatic stress and trauma- related disorders.
	Identify how traumatic experiences and their responses affect people with IDD.

LEARNING OBJECTIVES

Mo	dule	e 4:Well-Being & Resilience
		Explore the impact on caregivers of learning about the traumatic experience of the person with IDD they care for.
		Explain strategies for strengthening protective factors to enhance well-being, resilience, and recovery.
		Describe frameworks for promoting a healing and protective environment in order to create a safe and meaningful life for adults with IDD.
		Identify person-centered planning techniques to help adults with IDD realize their hopes and dreams.
	dule atm	e 5: IDD- & Trauma-Informed Services & nent
		Explain how to enhance protective factors of people with IDD & caregivers through appropriate traumainformed services & treatment.
		Discuss how to utilize adapted screening, assessment & planning tools to identify IDD- & trauma-informed needs of people with IDD.
		Discuss strategies for adapting the core components of trauma-informed treatments for people with IDD.
		Identify strategies for partnering with agencies & cross-system collaboration.
Mo	dule	e 6: Provider Self-Care
		Describe the difference between secondary traumatic stress, burnout and vicarious trauma.
		Identify how burnout develops among providers.
		Discuss potential sources, warning signs, and effects of secondary traumatic stress and organizational stress.
		Implement steps to stress reduction and self-care.

Module One

Setting the Stage









Acknowledgements

- Adapted from "The Road to Recovery Toolkit"
- Available for free on the National Child Traumatic Stress Network learning center
- https://www.nctsn.org/resources/roadrecovery-supporting-children-intellectualand-developmental-disabilities-who-have





Notes

Advisory Board Ruth Brown, PhD. Ananda Amstadter, PhD. Michael Southam Gerow, PhD - Virgin University Susan Ko, PhD - Science 37 Adam Kaul, MD - Regional Education Assessment Crains Services and Nabilisation Adam Kaul, MD - Regional Education Assessment Crains Services and Nabilisation Michael (Shell) Schwartz, MA - SufePlace, TX

What's new in this version?

- Focuses on the specific needs of adults with IDD, the contexts in which adults live, the impacts of trauma across the lifespan on adults, and assessment and treatments for adults with IDD
- Updated research findings on prevalence of trauma and stressrelated disorders according to the most recent diagnostic manuals used by psychologists (DSM-5 and DM-ID 2)







Notes


Terminology



- Providers anyone who works with or provides services to people with IDD.
- Intellectual and/or Developmental Disabilities (IDD) people who may have a broad array of conditions, beginning before adulthood and expected to last into adulthood, that result in significant impairment in adaptive functioning.
- Person-first preferred by most professions. Puts the person first and the disability second (e.g., "a person with Down syndrome" versus "a Down's person"). Some self-advocates prefer identity-first language ("I am a disabled-person").
- Behaviors a sometimes pejorative term describing the actions of a person with IDD when they are distressed, angry, frustrated, bored, etc.

Self Care

- · Step out and take a break.
- Talk to someone you trust.
- Do something relaxing.





Module one: Learning objectives

What Will I Learn Today?

- Define Trauma-Informed Care

- Peanize trauma-informed Care
 Realize that people with IDD are at-risk for traumatic experiences.
 Articulate that recover from traumatic experiences is possible.
 Establish the important role of parents/caregivers and the support network on the road to recovery.
- Describe frameworks that will be used to highlight key concepts throughout the training.



Notes

What does it mean to be "Trauma Informed"?



• Realizes the widespread impact of trauma

SAMHSA's Four R's of Trauma-Informed Care



- Recognizes the signs and symptoms of trauma
- Responds by changing behaviors, policies, and services to take trauma into account



· Resists re-traumatization of clients and staff with trauma histories

Source: store.samhsa.gov TIP 57: Trauma-Informed Care in Behavioral Health Services

Three Levels of Trauma Services

- Trauma-Informed
 Universal intervention
 Focuses on empowerment, safety, choice
- Trauma-Sensitive

 - Secondary intervention
 Aims to address trauma through assessment and modification to services
- Trauma-Specific

 - Tertiary intervention
 Use of evidence-based, or promising, trauma treatments

The Institute on Trauma and Trauma-Informed Care (2019). Trauma-Informed Organizational Change Manual. University of Buffalo School of Social Work. Retrieved from: www.socialwork.buffalo.edu/ittic





Realize

Trauma & IDD: Scope of the Problem

Trauma & IDD: Scope of the Problem



@VCU

WITH

Trauma & IDD: Scope of the Problem



⊚VCU *⊌WITH*

NPR Special Series: Abused and Betrayed. The Sexual Assault Epidemic No One Talks About



https://www.npr.org/series/575502633/abused-and-betrayed

Notes			
			

Trauma & IDD: Scope of the **Problem**

• "... those numbers from household surveys don't include people living in institutions where, Harrell said, research shows people are even more vulnerable to assault. Also not counted are the 373,000 people living in group homes."

The Sexual Assault Epidemic No One Talks About : NPR. https://www.npr.org/2018/01/08/570224090.the Sexual-assault-epidemic-np-one-talks-about







Notes

History of Institutional Care

- Willowbrook State School and others
- Overcrowding
- Abuse
- Neglect
- Unethical medical experimentation







Trauma & IDD: Scope of the Problem

- Study in the United Kingdom found that approximately 1,400 reports of sexual abuse committed against people with intellectual disability are reported annually
 - · Only 6% reach the court
 - Only 1% result in conviction







16

Trauma & IDD: Barriers to Recovery

- High rates of out-of-home placements
- Providers often feel insufficiently equipped to support people with IDD and trauma histories.
- Treatments for people with IDD often focus on behavioral compliance.
- Traumatic stress can lead to changes in learning, behavior & physiology—may place people at risk for further trauma.





"We went to them and they had no idea how to help us."




®VCU ØWITH


Myths about People with IDD

- ital health treatme th IDD
- le with tual disa
- rking with this
- bility
- with it are protected from trau ntal age (i.e., babies); they do p member
- Vou everything you ne



Notes








Notes

The Ecological Model Community Provides Family Individual







Notes				
 · · · · · · · · · · · · · · · · · · ·				

MAKING THE CONNECTION: WHY AM I DOING THIS WORK?

Think about what brought you to this training today. Is the reason professional? Personal? Is there a person with intellectual and developmental disability, in particular, whom you are thinking about?

What is it that connects you to this work?

What would you like to know?

Essential Messages of Working with People with IDD who have **Experienced Trauma**

- Know that there's hope; recovery from traumatic experiences is possible.
- Recognize that a person with IDD may have had a traumatic experience(s), which can have profound effects.







What Can a Provider Do?

- Create a learning climate/culture—honor the person's, caregiver's and provider's experience.
- Identify any fears or biases I have about working with people with IDD who have had traumatic experiences.
- Identify ways to increase my knowledge & skills





Notes

ACTION PLANNING ESSENTIAL MESSAGES 1 & 2

Think about the person you identified at the end of Module I that connects you to this work. What can help that person and other adults with IDD who have experienced trauma? (Put an "X" in up to 3 boxes next to the ideas you think you would MOST like to emphasize in your daily practice.)

-	ESSENTIAL MESSAGE I
4	Know that there's hope; recovery from traumatic experiences is possible.
	Over the next 3 months, I will identify my personal fears and/or biases of working with adults with IDD. I will begin to utilize the resources described in this Toolkit to help me to address those fears with knowledge and skills training.
	Over the next 3 months, I will read at least 2 articles online, or in peer-reviewed journals that describe the prevalence of trauma among adults with IDD.
	Over the next 3 months, I will articulate that recovery from traumatic experiences for adults with IDD is possible, to at least 3 adults with IDD and their caregivers.
	ESSENTIAL MESSAGE 2
2	Recognize that adults with IDD may have traumatic experiences, which can have profound effects.
	Over the next 3 months, I will identify 3 ways in which traumatic experiences are impacting the quality of life for at least three adults with IDD. I will review these areas each time I meet with them.
	Over the next 3 months, I will identify concerns and hopes , of caregivers of 3 adults with IDD who have experienced trauma. I will review these areas each time I meet with them.
	(Write my own.) Over the next 3 months, I will

Recovery is Possible



Most importantly, provide HOPE that people with IDD can recover from trauma.


- Help (re-)establish a sense of safety
- Listen, hear & validate what happened to them
- Support the their caregiving system
- Provide trauma-informed support & services
- Provide trauma-informed treatment

™VCU ™ WITI	@ ∨	CU	4	<u>⊌</u> WI	T
--------------------	------------	----	---	-------------	---

INotes	
	

Module Two

Development, IDD, and Trauma



Module Two: Learning Objectives What Will I Learn Today?

- Define IDD and various types of disabilities.
- 2. Discuss developmental tasks across typical development.
- Describe how IDD and traumatic experiences may disrupt typical development.
- Identify the areas in which a person's functioning is affected by IDD & subsequent adaptations that must be made.





What is Development?

Development is the process of change by which people become able to handle even more complex levels of moving, thinking, feeling and relating to others.



Notes

_	,
	6



Types of Development

- Language
- Motor
- Cognitive
- Social/Emotional
- Adaptive

How does a person develop to his/her full potential?





Development

Language development / Communication

Expressive language







Development

Gross Motor

- Large muscle groups
 Walking, running

Fine motor development

- Precision movements
- Tying shoes, writing







Notes					

Development

Cognitive Development

- Attention
- Memory
- Spatial Ordering
- Sequential Ordering
- Problem Solving
- Social Thinking







Development

Social/Emotional

- Self identity
- Relationships
- Social norms
- Sensory preferences and sensitivities
- Emotion regulation







Development

ability to adjust to various demands of normal daily living:

Self-Care

activities such as eating, toileting, bathing

• interest in activities outside the home

* taking care of/assisting with household tasks & taking care of personal possessions
 * Health & Safety
 * knowledge of basic health activities & physical dangers







Notes					

Federal Definition of Intellectual and Development Disabilities

- Impacts available services for people & families
- Emphasizes functional limitations in adaptive behaviors:

 - Receptive & Expressive Language
 - Learning
 - Mobility
- Self-Direction
- Capacity for Independent Living
- Economic Self-Sufficiency
- Impacts how or what parts of the system(s) families need to know how to navigate



Traits vs. Skills

- Traits
 - A distinguishing characteristic or quality, especially of one's personal
 - An inherited feature or characteristic.
- Skills
 - Proficiency, facility, or dexterity that is acquired or developed through training or experience.
 - A developed talent or ability.









Types of Disability Genetic Neuromotor Mental health (e.g., ADHD, depression, anxiety) Neurological (e.g., autism, epilepsy) Intellectual Sensory (e.g., vision, hearing, sensory-integration)








Notes					







Notes				





Notes

Living with IDD

If you are participating in an in-person workshop, you may complete this activity as a group. You can use this page to write down any insights that you want to keep in mind in your own work.

If you are completing this course independently, think about how living with IDD effects the quality of life for the person and the family. Imagine people with various disabilities. What are some ways their daily functioning is impacted in these areas? Think about the challenges that come with each of these areas and start to think about how you, as a provider, can help support people with IDD.

Self Care	Communication - unable to ask for help	Mobility	Social and Romantic Relationships
Home Living	Community Use	Health	Safety
	Leisure	Education	

Living with IDD: A Family's Perspective

- Safety & health are ongoing concerns
- Adults with IDD may require extended parental and family involvement in care
- There may be increased stress in the home which leads to conflict between parents, siblings & other relationships
- Questions about the extent to engage in "disability culture"
- Concerns about how to help their person with IDD achieve a valued & happy life



- Development can still progress in adults with IDD; however, it may occur at a different pace.
- Traumatic experiences at any age and stage of development can interfere with a person's developmental accomplishments.
- When people endure multiple traumatic events over long periods of time they are especially likely to have difficulty meeting developmental tasks.



Traumatic Experiences & IDD Can Disrupt Development









Erikson's Theory of Development



Typical Age	Task	Existential Question
0 - 18 m	Trust	Can I trust the world?
2 - 4 yrs	Autonomy	Is it okay to be me?
4 - 5 yrs	Initiative	Is it okay for me to do, move, and act?
5 - 12 yrs	Industry	Can I make it in the world?
13 - 19 yrs	Identity	Who am I? Who can I be?
20 - 39 yrs	Intimacy	Can I love?
40 - 64 yrs	Generativity	Can I make my life count?
65 - death	Ego Integrity	Is it okay to have been me?
	0 - 18 m 2 - 4 yrs 4 - 5 yrs 5 - 12 yrs 13 - 19 yrs 20 - 39 yrs 40 - 64 yrs	0 - 18 m Trust 2 - 4 yrs Autonomy 4 - 5 yrs Initiative 5 - 12 yrs Industry 13 - 19 yrs Identity 20 - 39 yrs Intimacy 40 - 64 yrs Generativity



Notes

IDD & Trauma in Early Childhood



Attachment to primary
caregiver

- Development of visual & auditory perception
- Recognition of & response to emotional
- Develop greater independence & capacity to

- Changes in eating & sleeping
- Become passive, quiet
- Heightened startle response
- · Confusion about what's dangerous & who to go to for protection
- Fear of being separated from familiar people/places
- Engage in regressive behaviors





IDD & Trauma in Middle Childhood



- Manage fears, anxieties, & aggression
- Sustain attention for learning & problem solving
- Control impulses & manage physical responses to danger

- Unwanted & intrusive thoughts/images
- Preoccupation with moments from the traumatic experience
- Replay the traumatic event
- Develop intense, specific new fears linking to the original danger
- Oscillate between being avoidant & reckless
- Sleep or concentration challenges





IDD & Trauma in Adolescence and **Young Adulthood**



Developmental Tasks

- Think abstractly
- Anticipate & consider the consequences of behavior
- Accurately judge danger & safety
- Increased impulse control & ability to defer gratification by thinking through

Trauma's Impact

- May feel embarrassed or angry about bouts of fear & exaggerated physical responses
- Difficulty imagining or planning for the future; decreased motivation for learning
- Low self-esteem & helplessness
- Difficulties with trust
- Reckless &/or self-destructive behaviors (e.g., drugs, cutting)



Notes

IDD & Trauma in Adulthood



Developmental Tasks

- Dialectical thought
- Intimate Relationships and parenting
- Social Inclusion
- Work and career
- Managing declining physical functioning/medical concerns

Trauma's Impact

- Difficulty imagining or planning for the future;
- Decreased motivation for learning or working
- Difficulties with trust
- Isolation and social withdrawal



IDD & Trauma in Older Age



Developmental Tasks

- Continue living independently as long as possible
- Managing energy and physical changes
- Changes in health, cognition, and
- Generativity
- Making sense of one's life

Trauma's Impact

- Late-Onset Stress Symptomology (LOSS)
- Changes in cognition and memory
- Symptoms of trauma may be mistaken for delirium or dementia
- Decreased independence
- Difficulty making meaning of trauma in context of life







"Adults with Developmental Disabilities" https://youtu.be/MGIEpswIE5w



		 	_
 	 	 	_
 	 	 	_
 		 	_
 	 	 	_
 	 	 	_
 	 	 	_
 	 	 	_
 	 	 	_
 	 	 	_



Notes	

CASE VIGNETTE JACQUELINE

Jacqueline is a 34-year-old woman who grew up in the lower valley of El Paso, Texas. Jacqueline was diagnosed with Down's syndrome (trisomy 21) at birth and was identified with hearing impairment and a congenital heart defect early in her life. She was diagnosed with an intellectual disability in elementary school.



Jacqueline's parents divorced when she entered Kindergarten. She, her mother and two older sisters had

no option but to move in and live with extended family members. They finally settled in and lived with Jacqueline's mother's sister. Jacqueline's mother was working two minimum wage jobs to save up for a place to live. She also contributed to her sister's household funds and her children's need. Jacqueline was often left in the care of her aunt, her older sister or an older cousin still living at home.

There were times during her childhood when occupational, speech therapy and doctor appointments were missed. Jacqueline's older cousin Robert, tended to play rough with her, sometimes leaving bruises and he frequently made fun of her speech. Jacqueline also cried a lot and had frequent tantrums at school and home. At those times, her aunt's boyfriend would send her to her room after yelling at her to stop crying and being such a bad girl.

Her mother eventually remarried and at the age of eighteen Jacqueline was placed in a group home.

In the group home where Jacqueline lives with 3 other women with disabilities, she began to experience waking in the middle of the night from nightmares. In response, sometimes evening staff would go into her room and just be with her until she was able to go back to sleep.

Jacqueline's sister visits every weekend and recently noticed that Jacqueline is very defensive to the touch and became physically aggressive when she tried to give her a hug. Jacqueline's sister also noticed that the sheets and bedding were soiled and had not been changed. Staff said that she has started wetting the bed again.

Jacqueline's sister is concerned with the aggressiveness and the care that Jacqueline may not be receiving by the staff in the group home.

CASE VIGNETTE AUSTIN

Austin is a 22-year-old male who was a healthy baby and developed normally for the first year of his life. At 18 months of age, his mother started to notice that Austin didn't seem interested in or to have curiosity about other children. He began playing with only a few select toys. Shortly before the birth of his younger brother, Austin stopped making eye contact and by the time he was 21 months old, he had ceased speaking altogether (or making any sounds except when he cried). He seemed to be



withdrawing more and more into his own world. Progressively, loud sounds were distressing and he began plugging his ears, eventually loud noises that never used to bother him—became unbearable. Along with his auditory hyper-sensitivity came sensory issues in all areas. When he was 2 I/2, he began showing signs of "visual stimming" (repetitive visual stimulation) and very pronounced obsessive compulsive behaviors. Austin was formally diagnosed with autism spectrum disorder by a pediatric neurologist.

Austin grew up in a bilingual household. His mother speaks English to him and his father speaks Hungarian. At the age of three, he was found to be farsighted and given glasses. At 3 1/2 years he was placed in a special education preschool. He was entirely nonverbal by that time. Over the summer he attended an Early Education Center. Austin received private speech therapy through Easter Seals in addition to public school and began speaking after the weekly speech therapy visits. His speech was limited to "no," "bye," and "hi." He would repeat words if asked and also use single words to request familiar things. Occupational therapy was provided intermittently throughout this time. These interventions became a regular part of his life throughout his early childhood and elementary years.

At the age of I2-years-old, Austin's parents stopped his therapies and interventions. They decided he needed a break and felt he had made great progress and was functioning well at school and home. His parents wanted Austin to be in general education for at least some of his classes. In middle school he began to run away from the school and school security would call his parents to come and pick him up once they had restrained and isolated him. Austin began to show more aggression as he entered high school at home and school.

In high school and through his transition age school years Austin began to see a psychotherapist on a weekly basis to help him with managing his emotions and interactions with peers. Austin was diagnosed with Bi-Polar disorder when he was evaluated by a psychiatrist his senior year of school. As Austin transitions out of high school at 22-years-old, his parents would like him to live on his own and are working towards a plan for him to live close to them in his own apartment. Austin has had difficulty making and keeping friends and his parents hope by living on his own it will help him achieve some social connections.

CASE VIGNETTE JOEY

Joey, a 30-year-old with an intellectual disability from Wichita, KS was taken to have his blood drawn by his mother who typically accompanies him to medical visits a few years ago. The phlebotomist greeted Joey and his mother and escorted Joey to the chair for the blood draw. He then informed Joey's mother that she needed to wait in the waiting room. Joey's mother began to explain that she always accompanied Joey during his medical procedures, but the phlebotomist did not let her finish her explanation.



He stated that all parents are required to wait in the waiting room and he shut the door as she stepped out.

Joey then got out of the chair and began to pace around the room. The phlebotomist asked him to sit down, but Joey said he wanted to leave. The phlebotomist said that they would leave as soon as he sat down and completed the procedure. Joey hesitantly sat down, but was squirming in his seat and fidgety with his hands. The phlebotomist insisted with an angry tone that he hold still. Frightened, Joey began to leave his seat. The phlebotomist blocked him and grabbed his arm and held it down. He said to Joey "if you don't hold still, you're not going to be able to leave." Joey became increasingly fearful and began to yell and cry for his mother. The phlebotomist proceeded to continue the blood draw restraining his arm and body.

Now, Joey becomes upset whenever he has to go to a doctor's appointment. He cries and tries to hide and sometimes becomes aggressive. As a result, he has been missing regular medical appointments and his mother is worried that his health is declining.

CASE VIGNETTE KATIE

Katie is 35-year-old and has been working at her job in an office for almost a year. She has done well organizing supplies and keeping the conference rooms prepared for meetings. She gets along well with other employees and typically likes going to work each day. She is assisted by a job coach that supports her to stay focused on her tasks. While she sometimes gets distracted, she is congenial with all those supporting her.



Katie's job coach recently went on maternity leave for several months. The new job coach was a kind young man who was committed to helping Katie. Soon after he started the job, Katie became agitated on the job, wasn't relating well to other employees, and wasn't performing her tasks as was expected of her. Anytime the new job coach approached her to assist, Katie got angry and told him to go away.

Katie's employer noticed the changes and suggested that they consult with Katie's case manager to gain insight into what may have generated the change in Katie's work experience. It was discovered that in high school Katie had been sexually assaulted by a young man of similar size and build as the new job coach. Recognizing that Katie didn't feel safe (even though she was safe), enabled those supporting her to get her the support she needed to work through her fears.

CASE VIGNETTE TREVELYAN

Trevelyan is a 65-year-old man from Detroit, MI. He has a bright smile and loves physical activity. Trevelyan has autism and lives in a residential center for people with developmental disabilities and has since he was 13 years old when his mother felt that she could no longer care for him at home due to his size and aggressive behaviors. Trevelyan does not use verbal communication. He knows a few words in sign language ("please," "thank




you," and "sorry"). After bringing him to the residential center as a child, his mother and other family members have rarely visited him, stating that they couldn't afford to take time off work or travel the 1.5 hours to the center to visit him.

His case file indicates that he grew up in an impoverished area and that he likely had limited access to appropriate interventions, such as applied behavior analysis, speech and language therapy, or occupational therapy. Not much is known about his home life prior to coming to the center. Because he frequently exhibits aggressive behaviors towards staff and peers, he is often on one-to-one supervision. When he was younger, restraints and basket holds were frequently used in response to his aggressive behaviors.

A behavioral analysis indicated that Trevelyan is more likely to act aggressively when he isn't allowed to have a preferred snack (either it is unavailable or belongs to another resident), when another resident is doing something that he wants to do, or to escape demanding tasks at his day program. His behavior plan has focused on redirecting him to other tasks and minimizing task demands with limited success. However, there are other times when he will push staff, run out of the building, or start slamming into doors and walls with no apparent antecedent. It is assumed that he becomes "overstimulated" because of his autism. Staff are frequently injured and turnover has been high, and many staff report that the stress and injuries from working with Trevelyan is a major reason they quit.

PUTTING IT INTO PRACTICE CASE VIGNETTES: TRAUMA, IDD & DEVELOPMENT

	ead the case vignette your group has been assigned and identify the ways in which auma and intellectual and developmental disability are/may disrupt development.
•	Identify any behaviors and responses related to intellectual and developmental disability.
-	Identify any behaviors and responses related to traumatic experiences.
	What are other ways that intellectual and developmental disability and traumatic experiences may disrupt typical development for the age range (e.g., emerging adult adult, older adult) of the case you've been assigned?





What Can a Provider Do?

- Ask questions and gather information (formally or informally) to understand the person's developmental age & how IDD is affecting his/her functioning.
- Identify IDD-related support needs of the person & family.
- Identify how living with IDD is affecting the quality of the person's life.







ACTION PLANNING ESSENTIAL MESSAGES 3 & 4

Think about the person you identified at the end of Module I that connects you to this work. What can help that person and other adults with IDD who have experienced trauma? (Put an "X" in up to 3 boxes next to the ideas you think you would MOST like to emphasize in your daily practice.)

ESSENTIAL MESSAGE 3 Recognize a person's developmental level and how IDD and traumatic experiences are affecting his/her functioning. Over the next 3 months, I will identify ways in which **trauma and IDD** have disrupted development for at least 3 adults with IDD. I will review these areas each time I meet with these them. Over the next 3 months, I will identify at least one alternative way of communicating/teaching a therapeutic skill (e.g. deep breathing, muscle relaxation) to use in place of, or in addition to, verbal communication for at least 3 adults with IDD. I will practice the chosen skill with them each time we meet. Over the next 3 months, I will review my **bibliotherapy resources** to determine if I have enough books to share with adults at different developmental levels about at least 3 different types of trauma. **ESSENTIAL MESSAGE 4** Utilize a developmental lens when making meaning of a person's traumatic experiences and responses. Over the next 3 months, I will identify how **living with IDD** is affecting the quality of life in specific areas (e.g., self-care, communication, etc) for at least 3 adults with IDD who have had traumatic experiences. For each of these people, I will develop a plan, with the person and caregivers that addresses these specific areas. I will review this plan each time I meet with them during this 3-month Over the next 3 months, I will learn a new method of assessing the person's **intellectual abilities** that may be better-suited to working with an adult with IDD who has had a traumatic experience (e.g., learn an IQ test that is normed on a more diverse population, non-verbal methods, etc.).

(Write my own.) Over the next 3 months, I will...

Module Three

Traumatic Stress Responses in People with IDD



Module Three: Learning Objectives

What Will I Learn Today?

- 1. Identify the types of events that can cause traumatic stress.
- Define traumatic stress and trauma-related disorders.
- Identify how traumatic experiences


 3. and their responses affect people with IDD.



Recognize

the signs and symptoms of trauma





Notes



Types of Traumatic Experiences

- Single traumatic experience (e.g., serious car accident, disasters)
- Experiences that occur together (e.g., domestic violence & physical abuse)
- Experiences that can extend over time (e.g., sexual abuse)
- Mass trauma (e.g., natural disaster, civil unrest, pandemic)
- Experiences that are a mixture of the above









What is Traumatic Stress?

Personally <u>experiencing or witnessing</u> a real or perceived threat to emotional/physical well-being

Not everyone who experiences a traumatic event will develop symptoms of traumatic stress.

Ongoing <u>pattern of symptoms</u> may be diagnosed as PTSD or other mental health disorders










1 4000	
 	 —
 	 —
 	 —
 	
 	 —








		•
	 	•
	 	•
	 	•
 	 	•
		•

Traumatic Stress and Other Diagnoses

- Posttraumatic stress & grief reactions can develop into other mental health conditions & diagnoses over time (e.g., separation anxiety & depression)
- Exacerbate preexisting mental health problems
- Reactions to trauma reminders may sometimes result in misdiagnoses
- Competent assessment is critical for accurate diagnosis, and effective intervention





12 Core Concepts





Core Concept 1: Complexity

"There's a lot going on in the person's mind and body."

For a person with IDD

Moment-to-moment reactions can be even more complex due to physical, cognitive or communicative limitations that impact the range of protective actions that were possible during the traumatic experience.





Core Concept 2: Life Experience

"What the person brings to the experience from within and from the outside world effects the person's response."







Core Concept 2: Life Experience

"What the person brings to the experience from within and from the outside world effects the person's response."

For a person with IDD It is important to understand the traumatic experience within the context of how IDD affects the person's everyday life.





 	 _
 	 _
 	 _
 	 _
	_
 	 _
 	 _
 	 _
 	 _
 	 _
 	 _
 	 _
 	 _
 	 _
 	 _
 	 _
	_
	 _
	 _
	 _
	 _
	 _

Core Concept 3: Reminders & Adversities

"What's the world like that the person is living in afterwards?"



For a person with IDD
Trauma may add to significant secondary adversities related to IDD experienced by the person and family that existed prior to the traumatic experience.



Trauma Reminders

"For the longest time we couldn't understand why Billy would get so agitated for no apparent reason. We had no idea that the perfume his personal care attendant sometimes wore reminded him of his assault years ago."



Core Concept 4: Wide-Ranging Reactions

"A person's reaction to a traumatic event varies greatly."

trauma in adults with

For a person with IDD Recognizing reactions to IDD can be complicated by the wide-ranging expressions of their disabilities.



Notes	



Core Concept 5: Danger & Safety

"A person's ability to believe in and trust in a protective shield has been broken and must be restored."

For a person with IDD

Restoring trust requires both time and strategies which take the person's IDD into account.







Core Concept 7: Protective Factors

"Promoting and restoring."

For a person with IDD

Reassure the person and parents/caregivers that strategies for addressing the effects of traumatic experiences can be adapted for adults with IDD by enhancing protective factors.







Notes


Core Concept 8: Development

"Don't just think symptoms—think development."

For a person with IDD Traumatic experiences may result in a significant setback in developmental progress, which was already challenged by IDD.








Core Concept 10: Culture Culture Sayered in its meaning and layered in its effects.



People with IDD and their families are often part of many cultural identities and communities, including disability communities, which may impact the experience and expression and of trauma.







Core Concept 12: Provider Distress

"We are stewards not just of those who allow us into their lives, but of our own capacity to be helpful."

For a parent/caregiver or provider of an adult with IDD

Traumatic experiences may (re-)evoke concerns about the vulnerability of the person with IDD and their own inability to protect them from harm.



PUTTING IT INTO PRACTICE CASE VIGNETTES: RESPONSES TO TRAUMATIC **EXPERIENCES**

Read the case vignette your group has been assigned and identify details related to trauma and intellectual and developmental disability. Keep in mind that not every case

	ill touch on all of the Core Concepts we've just discussed. Refer to the 12 Core oncepts handout to remind you of the main point for each Core Concept.
•	Identify the traumatic experience(s) and any trauma/loss reminders.
	Identify the Core Concepts that are applicable to the case.
	Think about the person with intellectual and developmental disability that you identified in our first exercise on the Making the Connection handout—how do traumatic experiences impact this person? How would you apply the Core Concepts to this person?

12 Core Concepts

for Understanding Traumatic Stress in People with Intellectual and Developmental Disabilities

Core Concept 1: Complexity - "There's a lot going on in the person's mind and body."

Core Concept 2: Life Experience - "What the person brings to the experience from within and from the outside world effects their response."

Core Concept 3: Reminders & Adversities - "What's the world like that the person is living in afterwards?"

Core Concept 4: Wide-Ranging Reactions - "A person's reaction to traumatic events varies greatly."

Core Concept 5: Danger & Safety - "A person's ability to believe in and trust in a protective shield has been broken and must be restored."

Core Concept 6: Caregiving Systems - "Everyone is shaken up by what happened."

Core Concept 7: Protective Factors - "Promoting and restoring"

Core Concept 8: Development - "Don't just think symptoms—think development."

Core Concept 9: Neurobiology - "How do you know what's dangerous?"

Core Concept 10: Culture - "Culture is layered in its meaning and layered in its effects."

Core Concept II: Social Contract - "Protection has failed in some way."

Core Concept 12: Provider Distress - "We are stewards not just of those who allow us into their lives, but of our own capacity to be helpful."



Essential Messages 5. Recognize that in the aftermath of trauma, understanding traumatic stress responses is the first step in helping people regain their sense of safety, value and quality of life.

What Can a Provider Do?

- Learn about a person's moment-to-moment reactions during a traumatic event.
 Learn about their intrinsic and extrinsic factors that comprise the ecology within which the trauma occurred.
- Ask about trauma reminders.
- Ask how their perceptions of danger and safety may have changed following the traumatic experience.
- Explore how the cultural background of the person and caregivers may be influencing responses to the traumatic experience.
- Ask if they are involved with social institutions.







ACTION PLANNING ESSENTIAL MESSAGE 5

Think about the person you identified at the end of Module I that connects you to this work. What can help that person and other adults with IDD who have experienced trauma? (Put an "X" in up to 3 boxes next to the ideas you think you would MOST like to emphasize in your daily practice.)

5

ESSENTIAL MESSAGE 5

Recognize that in the aftermath of trauma, understanding traumatic stress responses is the first step in helping a person regain their sense of safety, value and quality of life.

Over the next 3 months, I will identify the moment-to-moment reactions people have during traumatic experiences with 3 adults with IDD.
Over the next 3 months, I will explore extrinsic factors (e.g., physical environment, community & cultural responses to trauma & IDD) for 3 adults with IDD who have experienced trauma.
Over the next 3 months, I will identify 2 trauma reminders for 3 adults with IDD who had experienced trauma. I will check in with them about their responses to these reminders each time I meet with them.
Over the next 3 months, I will identify the impact of the traumatic experience on the caregiving system for 3 adults with IDD who have experienced trauma.
Over the next 3 months, I will identify at least 3 protective factors that will help people to recover more quickly from the harmful effects of trauma or loss for 3 adults with IDD who had traumatic experiences. I will check in about these protective factors each time I meet with them.
Over the next 3 months, I will ask 3 parents/caregivers about their feelings about the traumatic experience of the person with IDD they support to assess whether they need a referral to address secondary traumatic stress .
(Write my own.) Over the next 3 months, I will

Module Four

Well-Being and Resilience





Module Four: Learning objectives

What Will I Learn Today?

- Describe frameworks for promoting a healing and protective environment in order to create a safe and meaningful life for adults with IDD. Identify person-centered planning techniques to help adults with IDD realize their hopes and



Notes

Self Care •Step out and take a break. •Talk to someone you trust. •Do something relaxing.





Notes					

Respond

Change behaviors, policies, and services to take trauma into account



Notes

Ask & Answer Questions

- Responses to Traumatic Experiences
- Cultural/Linguistic Factors
- Traumatic Experiences of Caregivers
- Family Dynamics
 - Current and historical dynamics
 - Needs of the Caregivers
- Existing Supports/Protective Factors







Provide Timely Support

- Prompt intervention, in response to traumatic experiences, can diminish the overall effects of traumatic stress in people with IDD using Psychological First Aid.
 - Stabilization
 - Information Gathering
 - Practical Assistance

 - Information on Coping
 - Linkage with Collaborative Services





All Behavior is Communication



"If You Listen, You Will Hear Us" https://youtu.be/Hp4PW17U_h8





Enhance Well-Being and Resilience

- Families and caregivers are a critical part of both protecting people from harm and enhancing their well-being.
- Providing trauma-informed education and services to parents and other caregivers enhances their protective capacities.
- Providers should recognize that caregivers themselves may have trauma histories or experience secondary traumatic stress.





Recognize Caregivers' Hopes & Concerns

Caregivers of people with IDD want:

- To find & provide the best care for the person they support
- To sustain & maximize the person's maturity and independence across development & changing circumstances
- To help them function at his/her best and attain a meaningful life



The Challenge for Caregivers

- Feelings of Inadequacy/ Not Knowing What to Do
- Grief & Loss
- Strain on Relationships
- Anticipatory Anxiety
- Secondary Traumatic Stress
- Stress due to ongoing needs of the person that has continued into adulthood






Address Traumatic Experiences of Caregivers

A personal history of trauma can:

- · Compromise caregivers' decision-making
- Interfere with their ability to form and maintain secure & trusting relationships (with the person with IDD, partners, & service providers)
- Impair caregivers' ability to regulate their emotions
- Lead to less effective coping strategies, including substance abuse
- Cause caregivers to be reminded of their own traumatic experiences





Promote Secure Attachment

- A highly responsive caregiving style is associated with a variety of benefits across developmental stages including into adulthood.

 - a Imagine the inner life of the person
 Create or re-create a nurturing/healing environment
 Provide comfort to help a person regulate
 Structure, routine & limit-setting
- Facilitate healthy relationships with peers in the community.


upport	8	100	0-	
******			S.III	100
- December 1	3	-	7	nation for
- friends				-

LOCAL RESOURCES

Begin this exercise by identifying 5 local organizations that you frequently refer people and their caregivers to or find helpful. Once you've created your own list, walk around the room and ask others to share their list—find at least 5 new organizations to add to your list.

Partner with Caregivers to Create a Recovery Team





Putting It Into Practice





Activity-Putting It Into Practice







Resist

Take steps to resist re-traumatization of clients and staff with trauma histories



Promote Person-Centered Planning

- Develop a treatment and support plan using child-centered planning
 - Identify what is important to the person
 - Identify what others need to do to support what is important to the person
- Identify key individuals involved in person's life
- Foster collaborative relationships among those individuals



Notes

Positive Behavior Support

- Effectively used with individuals with IDD & mental health needs
- Different from behavior management
- Starts with the premise that individuals are coping in the most effective way they know
- Change the environment so that individuals can get what they want & teach skills so they have more effective tools to get what they want or cope when they can't









Gentle Teaching

- Developed by John McGee to interact with people with IDD & challenging behavior.
 Basic premise is that challenging behavior is a reflection of life experience often characterized by absence of quality of life &/or trauma.
- by assence of updany of me activities.

 Focuses on interacting in ways that improve quality of life for the person with IDD through meaningful & positive reciprocal relationships.

 Relationships are understood in GT as transacted in repetitive micro-behaviors using the tools of our hands, eyes, voice & presence.

https://gentleteaching.com/






Gentle Teaching



Notes	

Positive Identity Development

- Developed by Karyn Harvey, PhD to assist individuals with IDD in revealing their true potential as human beings & experiencing the happiness in life that they so deserve
- Positive identity development for people with IDD focuses on discovering, defining & celebrating the self vs taking on the sole identity of "disabled"
- Basic premise is that an approach modeled on psychological well-being vs ineffective behaviors leads to human health & authentic







Positive Identity Development

- Pleasure: things we enjoy in a predominantly happy state (e.g., eating, watching TV)
 - Passive
- Engagement: actively participating in something that is enjoyable (e.g., cooking, hiking)
 - Often leads to the development of skills & talents
- Meaning the act of making a real difference in the world of others through use of one's own strengths

http://www.pursuit-of-happiness.org/teaching-resources/







Positive Identity Development

Materials accompany Positive Identity Development: An Alternative Treatment Approach for Individuals with Mild and Moderate Intellectual Disabilities (Harvey, 2009)

Available at pid.thenadd.org

My Book About Recovery!

This is the book about Traveling the Long Road to Recovery Ву






Notes

Trauma-Informed Behavioral Interventions

- · Rich descriptions and case vignettes of effects of trauma on people with IDD.
- Strategies for trauma-informed crisis prevention and intervention.
- Examples of trauma-informed behavior support plans

https://www.aaidd.org/publications/bookstore-home/product-listing/trauma-informed-behavioral-interventions-what-works-and-what-doesnot



Notes

Essential Messages 6. Utilize an DD- & trauma-informed person-centered approach to support both the adult with IDL & the caregivers. 7. Help caregivers, and other professionals in the person's life, strengthen projective factors.

What Can a Provider Do?

- Identify and enhance strengths and natural supports.
- Ask questions about what they need.
- Take the time to listen to and ask them what about their questions, concerns & challenges.
- Provide practical tools for promoting a secure attachment and a healing & protective environment.
- Provide anticipatory guidance.
- · Help find access to respite care.







ACTION PLANNING ESSENTIAL MESSAGES 6 & 7

Think about the person you identified at the end of Module I that connects you to this work. What can help that person and other adults with IDD who have experienced trauma? (Put an "X" in up to 3 boxes next to the ideas you think you would MOST like to emphasize in your daily practice.)

ESSENTIAL MESSAGE 6 Utilize an IDD- & trauma-informed person-6 centered approach to support both the adult and the caregivers. Over the next 3 months, I will identify at least 3 activities that adults with IDD who have had traumatic experiences find meaningful. I will check in with them about these activities each time we meet. Over the next 3 months, I will identify at least 3 life goals (in their own words), that adults with IDD who have had traumatic experiences have for themselves. I will check in about these life goals each time we meet. Over the next 3 months, I will identify at least 3 adults with IDD who have siblings. I will work with the caregivers to identify needs of the siblings. **ESSENTIAL MESSAGE 7** Utilize a developmental lens when making meaning of a person's traumatic experiences and responses. Over the next 3 months, I will identify 3 **protective factors** to strengthen for 3 adults with IDD who had traumatic experiences. For each person, I will review these protective factors each time we meet during this 3-month period and develop strategies for enhancing them. Over the next 3 months, I will identify 3 adults with IDD, with whom I can work to **increase happiness** through activities that provide engagement and meaning. I will ask them about the activities each time we meet. Over the next 3 months, I will identify 3 adults with IDD to work with on building a recovery team. I will identify ways (in person, by phone, or by email) that I can help facilitate the participation of other providers on the recovery team.

(Write my own.) Over the next 3 months, I will...

Module Five

IDD- and Trauma-Informed Services and Treatment



Module Five: Learning objectives

What Will I Learn Today?

- Explain how to enhance protective factors of people with IDD & caregivers through appropriate trauma-informed services & treatment.
- Discuss how to utilize adapted screening, assessment & planning tools to identify IDD- & trauma-informed needs of people with IDD.

 Discuss strategies for adapting the core components of trauma-informed treatments for re
- being precise of data anionical deathers of data anionical solutions of data anionical



Multiple Providers, **Multiple Hats** Sometimes a provider has multiple roles. Most of the time, multiple providers each wear multiple hats.



Developing a Recovery Plan



Trauma-Related Needs



Identify IDD-Related Support Needs



- Developmental assessment & screening provide care providers with the opportunity to
 - Understand how the person is functioning
 - Understand how treatment may need to be modified to support the person
- Input from caregivers is invaluable in understanding the person's functioning in relation to his/her peers





7
/

Identify IDD-Related Support Needs



- World Health Organization Disability Assessment Schedule 2.0
- Can be licensed for commercial or non-commercial use
- Available in many languages

 - Available in many languages
 3 versions
 36-lems across 6 domains of functioning
 Cognition
 Mobility
 Mobility
 Getting Along with Others
 Life Activities
 Participation in Community Activities
 12-lem brid assessment of overall functioning
 12-4 version

- 3 Formats
 Interviewer-administered
 Self-report
 Proxy-administered (e.g. parent/caregiver)

Source: http://www.who.int/classifications/icf/whodasii/en/





Notes

Identify IDD-Related Support Needs



- The Arc's Self-Determination Scale Adult Version
- Right of people to have control in their lives.
- Original Version
 - https://thearc.org/wp-content/uploads/forchapters/SD%20Scale%20Procedu ral%20Guidelines.pdf
- Adult Version
 - https://rtcom.umn.edu/database/instruments/asds





Identify IDD-Related Support Needs



Developmental Assessment

- A more in-depth exploration of the IDD and/or impact on development
- Venues for Assessment
- Regional Center/Mental Health System
 - Neuropsychological / psychoeducational assessment
- Medical System
- Legal System



	_		
	8		3

Identify IDD-Related Support Needs



- Examples of Assessment Tools

 - Intelligence assessments
 Wechsler Adult Intelligence Scales
 Stanford-Binet Intelligence Scale
 Woodcock-Johnson Tests of Cognitive Ability
 Adaptive functioning assessments
- Adaptive functioning assessments
 Violated
 Adaptive Behavior Assessment Systems
 Assessment of Functional Living Skills
 Developmental Assessment for Individuals with Severe Disabilities
 Non-verbal assessments
 Wechsier Nonverbal Scale
 Universal Nonverbal Intelligence Test
 Letter International Performance Scale
 Be aware of floor effects!
 May not accurately portray the relative strengths and weaknesses





Notes

Identify IDD-Related Support Needs



- People with IDD may have more or less facility with communication
- Communication may be a struggle because a person may be nonverbal &/or use assistive communication devices
- Complex & chronic medical comorbidities (e.g., chronic lung disease, seizures & complex feeding problems) may require reliance on technologies (e.g., tracheostomies, home oxygen & gastrostomy tubes)
- Intensive coordination of services & therapies may detract from trauma-focused and related behavioral needs
- Disability Care Coordination Organizations





Identify Trauma-Related Support Needs



- Not everyone needs trauma-focused treatment.
- Trauma screening and assessment help identify the people who most need trauma-focused therapy.
- Consider the type of trauma experienced when making service referrals & service plans.
- Assess the functioning of the caregiving system, including the impact of caregiver trauma.







- Involve the person with IDD to the extent possible.
 - May need to make use of an intermediary (e.g., sign language interpreter, professional familiar with assistive devices)
- Providers should pay attention to
 - Pace (e.g., slow down speech)
 - Complexity (e.g., use simple language)
 - Timing (e.g., present one concept at a time)
 - Sequencing (e.g., rearrange questions to build on strengths)
- Involve a wide range of caregivers in the assessment process (e.g., parents, siblings, paid support staff, day program staff)





Identify Trauma-Related Support Needs Screening



- Does not necessarily have to be administered by a mental health professional
- Usually includes questions regarding exposure to trauma & related symptoms
- Assists providers in understanding the person's & family's history & potential trauma reminders
- Directs trauma-informed case planning; positive screen may result in referral for comprehensive trauma mental health assessment





Identify Trauma-Related Support Needs



Examples of trauma screening tools for adults with IDD include

Bangor Life Events Schedule for Intellectual Disabilities (BLESID)

- ngor Life Events Schedule for Intellectual Disabilities (BLESID)
 Self-report and caregiver-report versions available at
 http://www.leehw.com/blesid/
 + Hulbert-Williams, L. Hastings, R. P. crowe, R., & Pemberton, J. (2011).
 Self-Reported Life Events, Social Support and Psychological Problems in
 Adults with intellectual Disabilities. Journal of Applied Research in
 Intellectual Disabilities, 24(5), 427–438.

 https://doi.org/10.1111/j.1468-1348.2011.00624.x

Lancaster and Northgate Trauma Scales (LANTS)

- nicaster and Northgate Irauma scales (LANIS)
 Self-report and caregiver-report versions
 Wigham, S., Hatton, C., & Taylor, J. L. (2011). The Lancaster and
 Northgate Trauma Scales (LANTS): The development and psychometric
 properties of a measure of trauma for people with mild to moderate
 intellectual disabilities. Research in Developmental Devaltities, 32(6),
 1651–1659. https://doi.org/10.1016/j.inid.2011.06.008





 	 	 -
 	 	 -



Examples of trauma screening tools for adults with IDD include

Impact of Events Scale - Intellectual Disabilities

- Self-report version
 - Hall, J. C., Jobson, L., & Langdon, P. E. (2014). Measuring symptoms of post-traumatic stress disorder in people with intellectual disabilities: The development and psychometric properties of the Impact of Event Scale Intellectual Disabilities (IES-IDs). The British Journal of Clinical Psychology, 53(3), 315–332. https://doi.org/10.1111/bjc.12048





Identify Trauma-Related Support Needs



- Many caregivers have histories of trauma (in childhood & adulthood).
- Trauma can impact caregiving & protective capacities.
- Awareness of caregiver trauma history helps providers better understand caregivers and link them to appropriate services.
- Examples of screening tools for caregivers:
- Parent Emotional Reaction Questionnaire
- Life Events Checklist
- Trauma Recovery Scale





Identify Trauma-Related Support Needs



Trauma Assessment

- A more in-depth exploration of the nature & severity of the traumatic events, the impact of those events, current trauma-related symptoms & functional impairment
- Includes a clinical interview, use of objective measures, behavioral observations of the person, and collateral contacts with family, caseworkers, etc
- Usually conducted by a mental health provider to drive treatment planning
- May occur over at least 2-3 sessions
- Should be used to determine the treatment plan







- Domains covered include:
 - Basic demographics
 - Family history
 - Trauma history (comprehensive, including events experienced or witnessed)
 - Developmental history
 - Overview of person's problems/symptoms





Identify Trauma-Related Support Needs









Classification Systems









Adaptations to Diagnostic Criteria for PTSD

As recommended by DM-ID 2



- B1. Distressing Memories: Behavioral acting out and self-injury may be symptomatic for those with greater cognitive impairment
- cognitive impairment Dreams: Distressing dreams without recognizable content may be more common in people with more severe cognitive impairment.

 B3. Dissociative Reactions: Trauma-specific reenactments may be confused with symptoms of psychosis in adults with IDD.
- B4. Distress to Internal or External Cues: May appear as disorganized or agitated behavior in people with greater cognitive impairments





Adaptations to Diagnostic Criteria for PTSD

As recommended by DM-ID 2

- C. Avoidance of Trauma Cues: Like trauma exposure threshold, the threshold for what may activate a trauma reaction may lower and broaden and result in a lowering and broadening of the avoidance
 - May be reported as "noncompliance" for individuals unable to verbalize reasons for avoidance
- D1. Inability to Recall Details of Traumatic Event: May be associated with cognitive impairment and requires careful assessment



Identify Trauma-Related Support Needs



DM-ID-2 Workbook

- Structured diagnostic method for mental health professionals.
- Incorporates structured caregiver interview for observable signs
- Guides clinicians in making more accurate diagnoses.







Identify Trauma-Related Support Needs



IDD- & Trauma-Informed Services & Treatment

- Trauma-Informed Services: refers to a broad array of support & care, including treatment
 - Advocacy
 - Case Coordination
- Trauma-Informed Treatment: refers to interventions that focus on addressing symptoms & responses to traumatic experiences
 - Continuous Assessment
 - Cross-System Partnerships





		 	_
		 	-
		 	-
		 	_
		 	_
		 	_
		 	_
		 	-
		 	_



Developing IDD- & Trauma-Informed Recovery Plans

- Identify what is important to the person.
 Recognize the person's talents and strengths as well as needs.
 Include the person's personal circle of significant relationships when developing a plan.
 Identify the person's hopes and dreams, as well as fears and nightmares.
 Identify additional information that you need to gather.
 Trauma assessment/screening
 Protective factors to enhance
 Focus on concrete action to move toward the desired life.
 Referrals to trauma-focused treatment
 Referrals to other supportive services (e.g., respite, activities, parent trauma)



	d	00	
		100	
100	W	11	ł

1 10 003	
	
	
	
·	
	
	
	

PUTTING IT INTO PRACTICE: FISH BOWL

In this activity, we will be using a role play to consider the perspective of people with IDD and/or their caregivers and providers when accessing services.

Let's put this into practice. Split up into groups of 4. Once you have your groups, please take your chairs and put them in a cross formation so a pair of them is facing one another and the other pair is facing on another. It will look like a +.

At your tables, assign a pair of chairs facing each other to be a provider and a person with IDD and/or their family member, and the other pair to be two observers.

Now take a seat and either choose a case vignette from your Participant Manual or think of a person with IDD your caseload to role play.

Once the activity starts the provider will begin asking the person with IDD and/or their family member questions to start the process of getting them closer to a IDD- & trauma-informed recovery plan. Think of everything we've already talked about.


- Do you want to get information on the IDD diagnosis, or maybe the traumatic experience?
- Do you bring up screening and assessment?
- Identify risk or protective factors?

While the role play is occurring the two observers will watch and listen. At any point during the role play, the observers have the opportunity to pause the role play and give the provider helpful, supportive, or constructive feedback. When the feedback is done the role play can continue.

When I say stop, everyone will freeze, each group will stand and rotate clockwise, changing the roles of each group member. When I say continue, the group will carry on where the last one left off. Observers will then be able to pause the game as seen fit.

This will go on until everyone has a chance to be the provider, person with IDD and/or their family member, and the observer.

Once everyone has gone through all 3 roles, we will have a large group report out to talk about how that felt, what worked well, what you would chance next time you are confronted with the same situation, etc.



Trauma-Informed **Evidence-Based Treatments**

There are many different evidence-based trauma-informed treatments. A trauma-informed mental health professional should be able to determine which treatment is most appropriate for a particular person and/or their caregivers.

- Cognitive Processing Therapy
- Cognitive Therapy
- Individual CBT with a Trauma Focus
- Prolonged Exposure
- Eye Movement Desensitization and Reprocessing (EMDR)
- · For fact sheets on treatments:
- http://www.istss.org/treating-trauma/new-istss-guidelines.aspx





How do trauma treatments work?



- Many effective trauma treatments rely on cognitive behavioral strategies:
 - Teach stress management & relaxation skills to cope with unpleasant feelings about the trauma
 - Utilize "exposure strategies" (i.e., talking about the traumatic experience at a speed that controls the level of distress for the person)
 - Create a trauma "narrative" or story of what happened in order to master painful feelings about the experience & resolve the impact of the experience
 - Explore potentially untrue ideas about what happened & why
 - · Change unhealthy/incorrect views
 - May involve parents or other caregivers

	140	les .	
			
			
			
			
			-
			
			
			
			
			

Adaptations to Trauma-Focused Treatment

- Increased psychoeducation
 Building an emotion vocabulary
 Differentiating between thoughts, feelings, and behaviors
- · Modify pace
- Frequent review of concepts
 Practice skills multiple times in session
- Generalization Training

 - Increased opportunities to practice out of session
 Practice in as many settings as possible
 Enlist caregivers and support









Trauma-Focused CBT Adaptations for People with IDD

- Be creative in the ways in which the narrative is recorded
 - Dictate responses
 - Use a tape recorder, video or still camera
 - Role-play, sing or dance
 - Use play
- Slow down pace
 - Allow additional time to share thoughts & feelings about traumatic experience
- Recognize that repetition is helpful for learning
 - A person with IDD may return repeatedly to inaccurate or unhelpful cognitions

Source: Brian Tallant, LPC, Aurora Mental Health Center



Trauma Services & Treatment Adaptations for People with IDD

- Be sure that all members of the caregiving team are using the same language to address the traumatic experience(s).
- Check for comprehension—don't assume that the material is too complex for the person to understand.
- Use multisensory interventions (e.g., non-verbal, play therapy) & tools to assist in learning.
- Allow more time for the client to learn the skills & use more repetition.
- Work explicitly on generalization of skills to other environments.







Learning more about CBT strategies for People with Intellectual Disabilities

- Book: "Cognitive Behaviour Therapy for People with Intellectual Disabilities: Thinking creatively" by Andrew Jahoda, Biza Stenfert Kroese, and Carol Pert. Published by Palgrave Macmillan
- Book: "The Emotion Regulation Skills System for Cognitively Challenged Clients: A DBT-Informed Approach" by Julie Brown. Published by Guilford Press
- Online Course: "Tailoring Trauma-Focused Cognitive Behavior Therapy for Children with IDD." The National Child Traumatic Stress Network's Learning
 Center https://learn.nctsn.org/enrol/index.php?id=566
- Online Course: "Using Cognitive Behavioral Therapy with Persons with IDD." Relias

Academy https://reliasacademy.com/rls/store/browse/productDetailSingleSku.jsp?&productId=c1110648















IDD- & Trauma-Informed Services & Treatment: Questions to Ask

- Do you provide services to people with IDD who have had traumatic experiences?
 How do you approach therapy with people & their caregivers who have had traumatic experiences (regardless of whether they indicate or request trauma-informed treatment)?
- Do you provide trauma-specific or trauma-informed therapy?
 If so, how do you determine whether the person needs trauma-informed therapy (e.g., screening/assessment tools)?
- How familiar are you with evidence-based treatment models designed & tested for treatment of person trauma-related symptoms?
- Can you describe the core components of your treatment approach?



Other Services That Enhance Resilience & Recovery

- Mentoring programs
- Sports, arts, recreational activities
- Community service
- Wraparound programs





Notes

Trauma & IDD: Create an Integrated System of Care



Strategies for Cross-System Collaboration



- Cross-train on the intersection of trauma and IDD
- Jointly develop protocols regarding trauma and collaborative services that promote resiliency
- Create multi-disciplinary teams
- Schedule regular team meetings
- Co-locate staff in clinics & community "hubs"
- Identify & utilize cross-system assessment tools
- Share knowledge of local resources & trauma-informed treatment options
- Integrate information-sharing systems



Notes	

PUTTING IT INTO PRACTICE: CASE VIGNETTES: ACCESSING SERVICES

Read the case vignette your group has been assigned and answer the following questions. You may want to refer to the Protective Factors handout to give you some ideas.

Are there protective factors that you want to keep in mind when identifying screening/assessment and/or treatment for this person and their caregivers? Are there protective factors that can be built upon in treatment? If you are unaware of any protective factors how would you cultivate some with this person and/or their caregivers?

What challenges/barriers/risk factors do you anticipate coming up for this person and their caregivers? Do you think the challenges/risk factors will impact their ability to access services?

What are three things you could do to advocate for this person and/or their caregivers to help them overcome these challenges?

PROTECTIVE FACTORS CHECKLIST

☐ Ask & answer questions of the person with IDD and their caregivers
☐ Provide timely information
☐ Address traumatic experiences of the person with IDD and their caregivers
☐ Promote secure attachment
☐ Promote a healing & protective environment
☐ Help access IDD- & trauma-informed support and services
\square Partner with the person and caregivers to create a recovery team
☐ Help navigate systems of care, including systems challenges
☐ Offer community resources
□ Connect people and families with supportive peers and mentors
□ Refer for formal support if needed
☐ Encourage connection with any spiritual communities
☐ Connect the person and caregivers with cultural community
☐ Build & encourage family cohesion
☐ Promote supportive interactions between the person with IDD and their caregivers
☐ Build social skills
☐ Address cognitive functioning
☐ Promote self-efficacy skills



Essential Messages

- Partner with agencies and systems to ensure earlier and more sustained access to services.
- Ensure that trauma-informed person-centered services, treatments and systems drive the recovery plan.



What Can a Provider Do?

- Identify the developmental level of the person in order to guide appropriate selection of trauma screening/assessment tools, services & treatments.
- Facilitate the person's self-determination
- Coordinate continuous care through communication with therapists and other providers on a regular basis.
- Participate in cross-training with other agencies and systems on issues related to trauma and trauma-informed care.
- Identify and interview local individuals/agencies







ACTION PLANNING ESSENTIAL MESSAGES 8 & 9

Think about the person you identified at the end of Module I that connects you to this work. What can help that person and other adults with IDD who have experienced trauma? (Put an "X" in up to 3 boxes next to the ideas you think you would MOST like to emphasize in your daily practice.)

8

ESSENTIAL MESSAGE 8

Partner with agencies and systems to ensure earlier and more sustained access to services.

Over the next 3 months, I will identify 3 local agencies that provide advocacy for adults with IDD. I will attempt to make contact with at least I person in each agency.
Over the next 3 months, I will identify 3 local organizations that provide intensive case management or care coordination services for adults with IDD. I will attempt to make contact with at least I person in each organization.
Over the next 3 months, I will identify 3 local clinics/providers that provide trauma-informed assessments for adults. I will attempt to make contact with at least I person in each clinic.
Over the next 3 months, I will identify 3 local clinics/providers that provide trauma-informed treatment for adults. I will attempt to make contact with at least I person in each clinic.
(Write my own.) Over the next 3 months, I will

ACTION PLANNING ESSENTIAL MESSAGES 8 & 9

	ESSENTIAL MESSAGE 9
9	Ensure that trauma-informed person-centered services, treatments, and systems drive the recovery plan.
	Over the next 3 months, I will review the standardization, reliability, and validity information for I of the trauma screening or assessment tools I use to determine the appropriateness of use for adults with IDD.
	Over the next 3 months, I will identify a trauma screening tool and use it with 3 adults with IDD who have experienced trauma.
	Over the next 3 months, I will identify a tool to assess parent/caregiver trauma and use it with 3 caregivers of adults with IDD who have experienced trauma.
	Over the next 3 months, I will identify 3 adaptations I can make to my treatment of choice for 3 adults with IDD who have experienced trauma.
	(Write my own.) Over the next 3 months, I will

Module Six

Provider Self-Care



Module Five: Learning objectives

What Will I Learn Today?

- Describe the difference between secondary

 1. traumatic stress, burnout and vicarious trauma.
- 2. Identify how burnout develops among providers
- Discuss potential sources, warning signs, and 3. effects of secondary traumatic stress and organizational stress.
- 4. Implement steps to stress reduction and self-





Secondary Traumatic Stress (STS) refers to the development of post-traumatic stress symptoms caused by at least one indirect exposure to traumatic material.

- Burnout refers to emotional exhaustion, depersonalization, and a reduced feeling of personal accomplishment. Burnout develops as a result of general occupational stress.
- <u>Vicarious trauma</u> refers to changes in the inner experience of the therapist, resulting from empathetic engagement with a traumatized client. It focuses less on trauma symptoms and more on the changes that occur following cumulative exposure to another person's traumatic experience(s).





Provider Burnout

- Caseload that requires intensive case coordination
 - Need for Advocacy
 - Need for Collateral Engagement
- Longer therapeutic process & treatment times
- Working in a stressed system
- Identifying ways to adapt screening, assessment & interventions takes time







Notes

STRESS WARNING SIGNS

Fill in the blanks below using the text in parentheses as your guide to learn more about your own stress warning signs.


I know that my stress level is beginning to	affect my relationships when
(Identify a behavior or action verb)	(Identify a behavior or action verb)
and(Identify a behavior or action verb)	·
Other people in my life can tell that I am s	tressed out when I look
	, or when I (Name an adjective that describes how you appear)
Sound like (Name an unpleasant sound or noun)	•
When I am feeling overwhelmed, staying o	Onnected is (Write an adjective that says how you feel)
	when my stress level is entify a verb)
elevated.	

What can we do? Preventing and responding to burnout, secondary traumatic stress, or vicarious trauma

Self-care is the ability to engage in helping others without sacrificing other important parts of one's life.







SELF-CARE OPTIONS

Below are some tips for using the time that you do have for self-care effectively. Carve out time each day to engage in self-care, even if you only have 2 minutes.

If you have			0 minutes
	Breathe Stretch Daydream Take your stress temperature Laugh Doodle Acknowledge one of your accomplishments Say no to a new responsibility Complement yourself Look out the window	000000000	Evaluate your day Write in a journal Call a friend Meditate Tidy your work area Assess your self-care Draw a picture Dance Listen to soothing sounds Surf the web Read a magazine
	Spend time with your pet Share a favorite joke	3	0 minutes
	minutes Listen to music Have a cleansing cry Chat with a co-worker Sing out loud Jot down dreams Step outside for fresh air Enjoy a snack or make a cup of		Get a massage Exercise Eat lunch with a co-worker Take a bubble bath Read non-work related literature Spend time in nature Go shopping Practice yoga Watch your favorite television
	coffee/tea		show

Source: Transforming the Pain: A Workbook on Vicarious Traumatization. Saakvitne, Pearlman & Staff of TSI/CAAP (Norton, 1996)









BALANCING YOUR SELF-CARE

This assessment tool provides an overview of effective strategies to maintain self-care. Look through the list and see if there are one or two items you can start doing tomorrow.

Phys	Eat regularly (e.g. breakfast, lunch and dinner) Eat healthy Exercise Get regular medical care for prevention Get medical care when needed Take time off when needed Get massages Dance, swim, walk, run, play sports, sing, or do some other physical activity that is fun Take time to be sexual with yourself, with a partner Get enough sleep Wear clothes you like Take vacations, day trips, or mini-vacations Make time away from telephones Other:	Spirit	Make time for reflection Spend time with nature Find a spiritual connection or community Be open to inspiration Cherish your optimism and hope Be aware of nonmaterial aspects of life Try at times not to be in charge or the expert Be open to not knowing Identify what is meaningful to you and notice its place in your life Meditate Pray Sing Spend time with children Have experiences of awe Contribute to causes in which you believe Read inspirational literature (talks, music, etc.)
	Make time for self-Care Make time for self-reflection Have your own personal psychotherapy Write in a journal Read literature that is unrelated to work Do something at which you are not expert or in charge Decrease stress in your life Let others know different aspects of you Notice your inner experiencelisten to your thoughts, judgments, beliefs, and feelings Practice receiving from others Be curious Say "no" to extra responsibilities sometimes Other:	Worl	kplace or Professional Self-Care Take a break during the workday (e.g. lunch) Take time to chat with co-workers Make quiet time to complete tasks Identify projects or tasks that are exciting and rewarding Set limits with your clients and colleagues Balance your case load so that no one day or part of a day is "too much" Arrange your work space so it is comfortable and comforting Get regular supervision or consultation Negotiate for your needs (benefits, pay raise) Have a peer support group Develop a non-trauma area of professional
	Spend time with others whose company you enjoy Stay in contact with important people in your life Give yourself affirmations, praise yourself Re-read favorite books, watch favorite movies Identify comforting activities, objects, people, relationships, places and seek them out Allow yourself to cry Find things that make you laugh Express your outrage in social action, letters and donations, marches, protests Other:	Balar	interest Other:







Connection





Organizational Stress

Organizations, like individuals, are living, complex, adaptive systems. They are vulnerable to stress, particularly chronic & repetitive stress.







Impact of Organizational Stress on Providers Increased absenteeism

- Impaired judgment
- Unwillingness to accept extra work
- Low motivation
- Lower productivity
- Poorer quality of work
- Greater staff friction
- Higher staff turnover



Steps to Organizational Stress Reduction

- Practice principles of safety & empowerment
- Share successes & shortcomings
- Ensure regular & reflective supervision sessions
- Support open communication
- Hold multidisciplinary case conferences



Steps to Organizational Stress Reduction • De-stigmatize providers' person

- De-stigmatize providers' personal reactions to the work & prioritize self-care
- Provide mentoring to new professionals
- Support continuing education
- Encourage training on traumainformed care at all levels
- Provide respite for providers



Wellness Activities

- Provide wellness activities
- Offer mindfulness exercises during lunch breaks
- Support staff celebrations





Trauma-Informed Organizational Change





Essential Messages 10. Practice ongoing self-care in order to increase effectiveness in delivering high quality support, services and treatment

What Can a Provider Do?

- Identify potential sources, warning signs and effects of STS and organizational stress.
- Identify strategies for stress reduction.
- Practice the ABC's of self-care.
- Identify wellness activities that could be provided in your organization.
- Identify self-care resources that you'd like to explore further.







Notes

ACTION PLANNING ESSENTIAL MESSAGE 10

Think about the person you identified at the end of Module I that connects you to this work. What can help that person and other adults with IDD who have experienced trauma? (Put an "X" in up to 3 boxes next to the ideas you think you would MOST like to emphasize in your daily practice.)

10

ESSENTIAL MESSAGE 10

Practice ongoing self-care in order to increase effectiveness in delivering high quality support, services and treatment.

Over the next 3 months, I will designate a time to consider whether I am noticing the warning signs of stress in my relationships. I will utilize a self-care strategy when I notice that I am experiencing stress.
Over the next 3 months, I will choose 3 tips for engaging in self-care from the Self-Care Options handout and practice I on a daily basis.
Over the next 3 months, I will choose 3 strategies for maintaining self-care that I identified on the Balancing Your Self-Care handout and practice I on a weekly basis.
(Write my own.) Over the next 3 months, I will

Personal Trauma-Informed Practice Action Plan

Go back to the strategies you identified under the Essential Messages at the end of each module. Then, select three strategies that you want to commit to implementing as part of your Personal Action Plan. Write each of these strategies in the boxes provided below, and in the corresponding box in the left hand column, write in the Essential Message number associated with each strategy.

ESSENTIAL MESSAGES

- I. Know that there's hope; recovery from traumatic experiences is possible, including for adults with IDD.
- 2. Recognize that a person with IDD may have had a traumatic experience(s), which can have profound effects.
- 3. Recognize a person's developmental level and how IDD and traumatic experiences are affecting his/her functioning.
- 4. Utilize a developmental lens when making meaning of a person's traumatic experiences & responses.
- 5. Recognize that in the aftermath of trauma, understanding traumatic stress responses is the first step in helping someone regain their sense of safety, value and quality of life.
- 6. Utilize an IDD- & trauma-informed person-centered approach to support both the person and their support network.
- 7. Help parents/caregivers, and other professionals in the person's life, strengthen protective factors.
- 8. Partner with agencies and systems to ensure earlier and more sustained access to services.
- 9. Ensure that trauma-informed person-centered services, treatments, and systems drive the recovery plan.
- 10. Practice ongoing self-care in order to increase effectiveness in delivering high quality support, services and treatment.

Personal Trauma-Informed Practice Action Plan

Essential Message	Strategies I will implement over the next 3 months (refer to the strategies you identified under Essential Messages at the end of each module)
Example:	
10	Over the next 3 months, I will set aside time to meet with each of my supervisees to assess risks for burnout or vicarious trauma and support their self-care.
	Strategy I
	Strategy 2
	Strategy 3

References

Abused And Betrayed. (n.d.). NPR.Org. Retrieved July 20, 2021, from https://www.npr.org/series/575502633/abused-and-betrayed

BLESID—Bangor Life Events Schedule for Intellectual Disabilities — Lee Hulbert-Williams, PhD. (n.d.). Retrieved July 20, 2021, from https://leehw.com/blesid/

Bruno Kohfield-Galeano. (n.d.). The Quality of Life (Documentary about Intellectual Disability) (2015). Retrieved August 13, 2021, from https://www.youtube.com/watch?v=XWNixFvKiaU&t=661s

Center for Advanced Studies in Child Welfare. (n.d.). Reflective Supervision & Consultation: Perspectives from Child Welfare Professionals. Retrieved August 13, 2021, from https://www.youtube.com/watch?v=opnneWili6M

CFScalesMeasures.pdf. (n.d.). Retrieved July 20, 2021, from http://www.psychink.com/rfiles/CFScalesMeasures.pdf

CONNECT -CST North. (n.d.). Trauma Informed Behavior Support Plans with Dr Karyn Harvey 2021. Retrieved August 13, 2021, from https://www.youtube.com/watch?v=xatOcqtfttl

Creative Options Regina. (n.d.-a). Gentle Teaching and Trauma Informed Support, a conversation with Karyn Harvey, Ph.D. Retrieved August 13, 2021, from https://www.youtube.com/watch?v=A6CTUUMz7zo

Creative Options Regina. (n.d.-b). *The Power of Gentle Teaching*. Retrieved August 13, 2021, from https://www.youtube.com/watch?v=NpW|KxcmXsl

Creative Options Regina. (n.d.-c). What is our Central Purpose in a Culture of Gentleness. Retrieved August 13, 2021, from https://www.youtube.com/watch?v=lwiPB]7ukvc

Dan Cleberg. (n.d.). People with Developmental Disabilities WORK. Retrieved July 20, 2021, from https://www.youtube.com/watch?v=LCa02lgV010

Davison, E. H., Pless, A. P., Gugliucci, M. R., King, L.A., King, D.W., Salgado, D. M., Spiro, A., & Bachrach, P. (2006). Late-Life Emergence of Early-Life Trauma: The Phenomenon of Late-Onset Stress Symptomatology Among Aging Combat Veterans. *Research on Aging*, 28(1), 84–114. https://doi.org/10.1177/0164027505281560

Disabled And Here. (n.d.). Retrieved July 20, 2021, from https://affecttheverb.com/disabledandhere/

FilmRise Documentaries. (n.d.). *Unforgotten:Twenty-Five Years After Willowbrook - Full Movie*. Retrieved August 13, 2021, from https://www.youtube.com/watch?v=HNLyKW8fCNg

Hall, J. C., Jobson, L., & Langdon, P. E. (2014). Measuring symptoms of post-traumatic stress disorder in people with intellectual disabilities: The development and psychometric properties of the Impact of Event Scale-Intellectual Disabilities (IES-IDs). British Journal of Clinical Psychology, 53(3), 315–332. https://doi.org/10.1111/bjc.12048

Home. (n.d.). Retrieved July 20, 2021, from https://www.aaidd.org/

Hulbert-Williams, L., Hastings, R. P., Crowe, R., & Pemberton, J. (2011). Self-Reported Life Events, Social Support and Psychological Problems in Adults with Intellectual Disabilities. *Journal of Applied Research in Intellectual Disabilities*, 24(5), 427–436. https://doi.org/10.1111/j.1468-3148.2011.00624.x

ISTSS - New ISTSS Prevention and Treatment Guidelines. (n.d.). Retrieved July 20, 2021, from https://istss.org/clinical-resources/treating-trauma/new-istss-prevention-and-treatment-guidelines

Keesler, J. M., Green, S.A., & Nochajski, T. H. (2017). Creating a Trauma-Informed Community Through University-Community Partnerships: An Institute Agenda. *Advances in Social Work*, 18(1), 39–52. https://doi.org/10.18060/21298

King, L.A., King, D.W., Vickers, K., Davison, E. H., & III, A. S. (2007). Assessing late-onset stress symptomatology among aging male combat veterans. *Aging & Mental Health*, 11(2), 175–191. https://doi.org/10.1080/13607860600844424

Lauer, E., & McCallion, P. (2015). Mortality of People with Intellectual and Developmental Disabilities from Select US State Disability Service Systems and Medical Claims Data. *Journal of Applied Research in Intellectual Disabilities*, 28(5), 394–405. https://doi.org/10.1111/jar.12191

lauralipsky. (n.d.). *Trauma Stewardship*. Retrieved July 20, 2021, from https://www.youtube.com/watch?v=tAKPgNZi as

Learn: Psychological First Aid (PFA) and Skills for Psychological Recovery (SPR). (n.d.). NCTSN Learning Center. Retrieved July 20, 2021, from https://learn.nctsn.org/course/index.php?categoryid=11

Leicestershire Partnership NHS Trust. (n.d.). If You Listen, You Will Hear Us. Retrieved July 20, 2021, from https://www.youtube.com/watch?v=Hp4PW17U h8

Life Events Checklist for DSM-5 (LEC-5) - PTSD: National Center for PTSD. (n.d.). [General Information]. Retrieved August 16, 2021, from https://www.ptsd.va.gov/professional/assessment/te-measures/life_events_checklist.asp

Materials for Positive Identity Development. (n.d.). *The NADD*. Retrieved July 20, 2021, from https://thenadd.org/materials-for-positive-identity-development/

National Child Traumatic Stress Network. (2018, March 5). The Road to Recovery: Supporting Children with Intellectual and Developmental Disabilities Who Have Experienced Trauma. The National Child Traumatic Stress Network.

https://www.nctsn.org/resources/road-recovery-supporting-children-intellectual-and-developmental-disabilities-who-have

Open Future Learning. (n.d.). ALL BEHAVIOR IS COMMUNICATION. Retrieved July 20, 2021, from https://www.youtube.com/watch?v=iZThnMP8N11

Pitman, R. K. (2010). Posttraumatic Stress Disorder and DementiaPosttraumatic Stress Disorder and Dementia. *JAMA*, 303(22), 2287–2288. https://doi.org/10.1001/jama.2010.767

Potter, C. M., Kaiser, A. P., King, L. A., King, D. W., Davison, E. H., Seligowski, A. V., Brady, C. B., & Spiro, A. (2013). Distinguishing late-onset stress symptomatology from posttraumatic stress disorder in older combat veterans. *Aging & Mental Health*, 17(2), 173–179. https://doi.org/10.1080/13607863.2012.717259

Prevalence and psychological sequelae of sexual abuse among individuals with an intellectual disability: A review of the recent literature—Gary Byrne, 2018. (n.d.). Retrieved July 20, 2021, from https://journals.sagepub.com/doi/10.1177/1744629517698844

Psychological First Aid. (n.d.). Coursera. Retrieved July 20, 2021, from https://www.coursera.org/learn/psychological-first-aid

Psychological First Aid Online Course | Red Cross. (n.d.). Retrieved July 20, 2021, from https://www.redcross.org/take-a-class/coronavirus-information/psychological-first-aid-online-course

Risk Reduction Book. (n.d.). Nora Baladerian, Ph.D. Retrieved July 20, 2021, from https://norabaladerian.com/risk-reduction-book/

RTC Media. (n.d.-a). Aging, Retirement, and End of Life for people with intellectual and developmental disabilities. Retrieved August 13, 2021, from https://www.youtube.com/watch?v=yA]O yGa9nA

RTC Media. (n.d.-b). Social Inclusion for people with intellectual and developmental disabilities. Retrieved July 20, 2021, from https://www.youtube.com/watch?v=-dDncNIVzwl

Shapiro, J. (2018a, January 8). The Sexual Assault Epidemic No One Talks About. In *All Things Considered*. NPR. https://www.npr.org/2018/01/08/570224090/the-sexual-assault-epidemic-no-one-talks-about

Shapiro, J. (2018b, January 20). In Their Own Words: People With Intellectual Disabilities Talk About Rape. *NPR*. https://www.npr.org/2018/01/20/577064075/in-their-own-words-people-with-intellectual-disabilities-talk-about-rape

ShawTVOkanagan. (n.d.). Adults with Developmental Disabilities. Retrieved July 20, 2021, from https://www.youtube.com/watch?v=MGIEpswIE5w

Teaching Resources | Pursuit of Happiness | Happiness is understandable, obtainable, and teachable. (n.d.). Retrieved July 20, 2021, from https://www.pursuit-of-happiness.org/teaching-resources-2/

The Arc's Self-Determination Scale, Adult Version (ASDS). (n.d.). Retrieved July 20, 2021, from https://rtcom.umn.edu/database/instruments/asds

The Institute on Trauma and Trauma-Informed Care (ITTIC). (n.d.). Retrieved July 20, 2021, from http://socialwork.buffalo.edu/ittic

TIP 57:Trauma-Informed Care in Behavioral Health Services | SAMHSA Publications and Digital Products. (n.d.). Retrieved July 20, 2021, from https://store.samhsa.gov/product/TIP-57-Trauma-Informed-Care-in-Behavioral-Health-Services/SMA14-4816

Trauma-Informed Organizational Change Manual. (n.d.). Retrieved July 20, 2021, from http://socialwork.buffalo.edu/social-research/institutes-centers/institute-on-trauma-and-trauma-informed-care/Trauma-Informed-Organizational-Change-Manual0.html

Wehmeyer, M. L. (n.d.). The Arc's Self-Determination Scale: 152.

WHO Disability Assessment Schedule (WHODAS 2.0). (n.d.). Retrieved July 20, 2021, from https://www.who.int/classifications/international-classification-of-functioning-disability-and-health/who-disability-assessment-schedule

Wigham, S., Hatton, C., & Taylor, J. L. (2011). The Lancaster and Northgate Trauma Scales (LANTS): The development and psychometric properties of a measure of trauma for people with mild to moderate intellectual disabilities. Research in Developmental Disabilities, 32(6), 2651–2659. https://doi.org/10.1016/j.ridd.2011.06.008

Yaffe, K., Vittinghoff, E., Lindquist, K., Barnes, D., Covinsky, K. E., Neylan, T., Kluse, M., & Marmar, C. (2010). Posttraumatic Stress Disorder and Risk of Dementia Among US Veterans. *Archives of General Psychiatry*, 67(6), 608. https://doi.org/10.1001/archgenpsychiatry.2010.61